

Diffractive ρ production with an AdS/QCD holographic wavefunction for the ρ meson

Jeff Forshaw* and Ruben Sandapen†

**University of Manchester, Oxford Road, Manchester M13 9PL, UK.*

†*Université de Moncton, Moncton, N-B, E1A 3E9, Canada
& Mount Allison University, Sackville, N-B, E4B 1E6, Canada.*

Abstract. We report on the results of our recent research published in [1] that shows that AdS/QCD generates predictions for the rate of diffractive ρ -meson electroproduction that are in agreement with data collected at the HERA electron-proton collider [2, 3]. Preliminary results of this research were presented in [4].

Keywords: Light-front holographic wavefunctions, AdS/QCD, diffractive ρ meson production

PACS: 11.25.Tq, 12.38.Aw, 12.38.-t, 13.60.Le

INTRODUCTION

We demonstrate another success of the AdS/QCD correspondence [5, 6, 7, 8] by showing that parameter-free AdS/QCD wavefunctions for the ρ meson lead to predictions for the rate of diffractive ρ meson production ($\gamma^* p \rightarrow \rho p$) that agree with the data collected at the HERA ep collider. To compute the rate for diffractive ρ production, we use the dipole model of high-energy scattering [9, 10, 11, 12] in which the scattering amplitude for diffractive ρ meson production is a convolution of the photon and vector meson $q\bar{q}$ light-front wavefunctions with the total cross-section to scatter a $q\bar{q}$ dipole off a proton. QED is used to determine the photon wavefunction and the dipole cross-section can be extracted from the precise data on the deep-inelastic structure function F_2 [13, 14]. This formalism can then be used to predict rates for vector meson production and diffractive DIS [15, 16] or to extract information on the ρ meson wavefunction using the HERA data on diffractive ρ production [17, 18]. Here we use it to predict the cross-sections for diffractive ρ production using an AdS/QCD holographic wavefunction for the ρ meson proposed by Brodsky and de Téramond [19]. With this AdS/QCD wavefunction, we also predict the second moment of the twist-2 Distribution Amplitude of the longitudinally polarized ρ meson. We find good agreement with Sum Rules and lattice predictions.

AN ADS/QCD HOLOGRAPHIC WAVEFUNCTION

The AdS/QCD wavefunction is given by [1, 19]

$$\phi(x, \zeta) = N \frac{\kappa}{\sqrt{\pi}} \sqrt{x(1-x)} \exp\left(-\frac{\kappa^2 \zeta^2}{2}\right) \exp\left(-\frac{m_f^2}{2\kappa^2 x(1-x)}\right) \quad (1)$$

where N is a normalization constant and m_f is the light quark mass.¹ In Eq. (1), $\zeta = \sqrt{x(1-x)}b$ where x is the light-front longitudinal momentum fraction of the quark and b is the transverse separation between the quark and antiquark. ζ is the variable that maps onto the coordinate z in the fifth dimension of AdS space [19]. The parameter $\kappa = M_\rho/\sqrt{2} = 0.55$ GeV [1].

This AdS/QCD wavefunction is rather similar to the original Boosted Gaussian (BG) wavefunction discussed in [20, 15]

$$\phi^{\text{BG}}(x, \zeta) \propto x(1-x) \exp\left(\frac{m_f^2 R^2}{2}\right) \exp\left(-\frac{m_f^2 R^2}{8x(1-x)}\right) \exp\left(-\frac{2\zeta^2}{R^2}\right). \quad (2)$$

If $R^2 = 4/\kappa^2$ then the two wavefunctions differ only by a factor of $\sqrt{x(1-x)}$, which is not surprising given that in both cases confinement is modelled by a harmonic oscillator [1]. In what follows we shall consider a parameterization that accommodates both the AdS/QCD and the BG wavefunctions:

$$\phi(x, \zeta) \propto [x(1-x)]^\beta \exp\left(-\frac{\kappa^2 \zeta^2}{2}\right) \exp\left(-\frac{m_f^2}{2\kappa^2 x(1-x)}\right). \quad (3)$$

The AdS/QCD wavefunction is obtained by fixing $\beta = 0.5$ and $\kappa = 0.55$ GeV where as the BG wavefunction is obtained by fixing $\beta = 1$ and treating κ as a free parameter.

The meson's light-front wavefunctions can be written in terms of the wavefunction $\phi(x, \zeta)$ [18]. For longitudinally polarized mesons,

$$\Psi_{h,\bar{h}}^L(b, x) = \frac{1}{2\sqrt{2}} \delta_{h,-\bar{h}} \left(1 + \frac{m_f^2 - \nabla^2}{M_\rho^2 x(1-x)}\right) \phi_L(x, \zeta), \quad (4)$$

where $\nabla^2 \equiv \frac{1}{b} \partial_b + \partial_b^2$ and h (\bar{h}) are the helicities of the quark (anti-quark) and for transversely polarized mesons,

$$\Psi_{h,\bar{h}}^{T=\pm}(b, x) = \pm [ie^{\pm i\theta} (x\delta_{h\pm, \bar{h}\mp} - (1-x)\delta_{h\mp, \bar{h}\pm}) \partial_b + m_f \delta_{h\pm, \bar{h}\pm}] \frac{\phi_T(x, \zeta)}{2x(1-x)}, \quad (5)$$

where $be^{i\theta}$ is the complex form of the transverse separation, \mathbf{b} . We impose the normalization condition:

$$\sum_{h,\bar{h}} \int d^2\mathbf{b} dx |\Psi_{h,\bar{h}}^\lambda(b, x)|^2 = 1, \quad (6)$$

where $\lambda = L, T$. This means we allow for a polarization dependent normalization (hence the subscripts on $\phi_{L,T}$).

¹ Here we shall take $m_f = 140$ MeV, which is the value used in the dipole fits to the deep-inelastic structure function, $F_2(x, Q^2)$ [18].




FIGURE 1. The χ^2 distribution in the (β, κ) parameter space. The AdS/QCD prediction is the white star.

COMPARISON TO DATA, SUM RULES AND THE LATTICE

We first compute the AdS/QCD prediction for the electronic decay width $\Gamma_{e^+e^-}$, which is related to the decay constant via

$$f_\rho = \left(\frac{3\Gamma_{e^+e^-}M_\rho}{4\pi\alpha_{\text{em}}^2} \right)^{1/2}.$$

Using

$$f_\rho = \frac{1}{2} \left(\frac{N_c}{\pi} \right)^{1/2} \int_0^1 dx \left(1 + \frac{m_f^2 - \nabla^2}{M_\rho^2 x(1-x)} \right) \phi_L(x, \zeta = 0), \quad (7)$$

we obtain $\Gamma_{e^+e^-} = 6.66$ keV, which is to be compared with the measured value $\Gamma_{e^+e^-} = 7.04 \pm 0.06$ keV [21].

To compute the rate for diffractive ρ production, we use the CGC[0.74] dipole model [13, 22] (see [18] for details), although the predictions do not vary much if we use other models that fit the HERA F_2 data [14, 23]. The plots comparing the AdS/QCD predictions and the HERA data can be found in [1]. Here, in figure 1, we show the χ^2 per data point² in the (β, κ) parameter space (see Eq. (3)). It confirms that the AdS/QCD prediction lies impressively close to the minimum in χ^2 . The best fit has a χ^2 per data point equal to 114/76 and is achieved with $\kappa = 0.56$ GeV and $\beta = 0.47$. Note that the BG prediction i.e. $\beta = 1, \forall \kappa$, is clearly further away from the minimum in χ^2 .

² We include the electroproduction data and also the decay constant f_ρ in the fit.

We have previously shown [18] that the twist-2 Distribution Amplitude (DA) can be related to $\phi_L(x, \zeta)$ according to

$$\varphi(x, \mu) = \left(\frac{N_c}{\pi}\right)^{1/2} \frac{1}{2f_\rho} \int db \mu J_1(\mu b) \left(1 + \frac{m_f^2 - \nabla^2}{M_\rho^2 x(1-x)}\right) \phi_L(x, \zeta) . \quad (8)$$

To compare to predictions using QCD Sum Rules [24] and from the lattice [25], we compute the moment:

$$\int_0^1 dx (2x-1)^2 \varphi(x, \mu) . \quad (9)$$

We obtain a value of 0.228 for the AdS/QCD wavefunction, which is to be compared with the Sum Rule result of 0.24 ± 0.02 at $\mu = 3$ GeV [24] and the lattice result of 0.24 ± 0.04 at $\mu = 2$ GeV [25]. The AdS/QCD wavefunction neglects the perturbatively known evolution with the scale μ and should be viewed as a parametrization of the DA at some low scale $\mu \sim 1$ GeV. Viewed as such, the agreement is good.

ACKNOWLEDGMENTS

RS thanks the organizing committee of Diffraction 2012 for their invitation, financial support and a very enjoyable conference.

REFERENCES

1. J. R. Forshaw, and R. Sandapen, *Phys.Rev.Lett.* **109**, 081601 (2012), 1203.6088.
2. S. Chekanov, et al., *PMC Phys.* **A1**, 6 (2007), 0708.1478.
3. The H1 Collaboration, *JHEP* **05**, 032 (2010), 0910.5831.
4. J. R. Forshaw, and R. Sandapen (2012), 1207.4358.
5. J. Erdmenger, N. Evans, I. Kirsch, and E. Threlfall, *Eur.Phys.J.* **A35**, 81–133 (2008), 0711.4467.
6. J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal, and U. A. Wiedemann (2011), 1101.0618.
7. M. S. Costa, and M. Djuric (2012), 1201.1307.
8. G. F. de Teramond, and S. J. Brodsky (2012), 1203.4025.
9. N. N. Nikolaev, and B. G. Zakharov, *Z. Phys.* **C49**, 607–618 (1991).
10. N. N. Nikolaev, and B. G. Zakharov, *Z. Phys.* **C53**, 331–346 (1992).
11. A. H. Mueller, *Nucl. Phys.* **B415**, 373–385 (1994).
12. A. H. Mueller, and B. Patel, *Nucl. Phys.* **B425**, 471–488 (1994), hep-ph/9403256.
13. G. Soyez, *Phys. Lett.* **B655**, 32–38 (2007), 0705.3672.
14. J. R. Forshaw, and G. Shaw, *JHEP* **12**, 052 (2004), hep-ph/0411337.
15. J. R. Forshaw, R. Sandapen, and G. Shaw, *Phys. Rev.* **D69**, 094013 (2004), hep-ph/0312172.
16. J. R. Forshaw, R. Sandapen, and G. Shaw, *JHEP* **11**, 025 (2006), hep-ph/0608161.
17. J. R. Forshaw, and R. Sandapen, *JHEP* **11**, 037 (2010), 1007.1990.
18. J. R. Forshaw, and R. Sandapen, *JHEP* **1110**, 093 (2011), 1104.4753.
19. G. F. de Teramond, and S. J. Brodsky, *Phys.Rev.Lett.* **102**, 081601 (2009), 0809.4899.
20. J. Nemchik, N. N. Nikolaev, E. Predazzi, and B. G. Zakharov, *Z. Phys.* **C75**, 71–87 (1997), hep-ph/9605231.
21. Particle Data Group, *J. Phys.* **G37**, 075021 (2010).
22. G. Watt, and H. Kowalski, *Phys. Rev.* **D78**, 014016 (2008), 0712.2670.
23. H. Kowalski, L. Motyka, and G. Watt, *Phys. Rev.* **D74**, 074016 (2006), hep-ph/0606272.
24. P. Ball, V. M. Braun, and A. Lenz, *JHEP* **08**, 090 (2007), 0707.1201.
25. P. A. Boyle, et al., *PoS LATTICE2008*, 165 (2008), 0810.1669.